

32. Jalife J, Berenfeld O, Skanes A, Mandapati R. Mechanisms of atrial fibrillation: mother rotors or multiple daughter wavelets, or both? *J Cardiovasc Electrophysiol* 1998;**9**:S2–S12.
33. Nademanee K, McKenzie J, Kosar E, Schwab M, Sunsaneewitayakul B, Vasavakul T, Khunnawat C, Ngarmukos T. A new approach for catheter ablation of atrial fibrillation: mapping of the electrophysiologic substrate. *J Am Coll Cardiol* 2004;**43**: 2044–2053.
34. Schauerte P, Scherlag BJ, Patterson E, Scherlag MA, Matsudaria K, Nakagawa H, Lazzara R, Jackman WM. Focal atrial fibrillation: experimental evidence for a pathophysiologic role of the autonomic nervous system. *J Cardiovasc Electrophysiol* 2001;**12**:592–599.
35. Roberts-Thomson KC, Stevenson IH, Kistler PM, Haqqani HM, Goldblatt JC, Sanders P, Kalman JM. Anatomically determined functional conduction delay in the posterior left atrium relationship to structural heart disease. *J Am Coll Cardiol* 2008;**51**:856–862.
36. Sauer WH, McKernan ML, Lin D, Gerstenfeld EP, Callans DJ, Marchlinski FE. Clinical predictors and outcomes associated with acute return of pulmonary vein conduction during pulmonary vein isolation for treatment of atrial fibrillation. *Heart Rhythm* 2006;**3**:1024–1028.
37. Rajappan K, Kistler PM, Earley MJ, Thomas G, Izquierdo M, Sporton SC, Schilling RJ. Acute and chronic pulmonary vein reconnection after atrial fibrillation ablation: a prospective characterization of anatomical sites. *Pacing Clin Electrophysiol* 2008;**31**:1598–1605.
38. Cheema A, Dong J, Dalal D, Marine JE, Henrikson CA, Spragg D, Cheng A, Nazarian S, Bilchick K, Sinha S, Scherr D, Almasry I, Halperin H, Berger R, Calkins H. Incidence and time course of early recovery of pulmonary vein conduction after catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2007;**18**:387–391.
39. Echahidi N, Pibarot P, O'Hara G, Mathieu P. Mechanisms, prevention, and treatment of atrial fibrillation after cardiac surgery. *J Am Coll Cardiol* 2008;**51**: 793–801.
40. Mason DP, Marsh DH, Alster JM, Murthy SC, McNeill AM, Budev MM, Mehta AC, Pettersson GB, Blackstone EH. Atrial fibrillation after lung transplantation: timing, risk factors, and treatment. *Ann Thorac Surg* 2007;**84**:1878–1884.
41. See VY, Roberts-Thomson KC, Stevenson WG, Camp PC, Koplan BA. Atrial arrhythmias after lung transplantation: epidemiology, mechanisms at electrophysiology study, and outcomes. *Circ Arrhythm Electrophysiol* 2009;**2**: 504–510.

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehq269

Online publish-ahead-of-print 30 July 2010

Computed angiogram of the upper extremities for diagnosing a rare cause of brachial arterial embolism: the 'Pitcher Syndrome'

Daniela Reutter¹, Roger Hunziker², and Marc Husmann^{1*}

¹Clinic for Angiology, University Hospital Zurich, 8091 Zurich, Switzerland; and ²Department of Radiology, University Hospital Zurich, Zurich, Switzerland

* Corresponding author. Tel: +41 44 255 3491/+41 44 255 2650, Fax: +41 44 255 4510, Email: marc.husmann@usz.ch

Embolic brachial artery occlusion most often originates from the heart, the aortic arch, or from subclavian artery aneurysms.

A 64-year-old man presented with acute upper limb ischaemia of the left arm. Duplex sonography confirmed a brachial artery occlusion, but no compression of the left subclavian artery or a subclavian aneurysm as source for the emboli. There were no signs for cardiac arrhythmia in the ECG nor cardiac thrombus by echocardiography. Computed angiogram of the thoracic and upper extremity revealed an aneurysm of the left posterior circumflex humeral artery (Panels A and B). Treatment consists in surgical embolectomy at the brachial bifurcation and in ligation of the aneurysm.

Aneurysms of the posterior circumflex humeral artery have been reported for volleyball and baseball players and named as the 'Pitcher Syndrome'. Repetitive traumas to the arterial wall during throwing motions of the shoulder affect the posterior circumflex humeral artery at the quadrangular space at the neck of the humerus. Embolic occlusion may result from dislocation of the squeezed thrombus from the aneurysmatic sack during shoulder movements. Since this patient did never practice any other sports than yoga, it might be assumed that certain specific yoga exercises with shoulder movements might have provoked the artery trauma.

The absence of cardiovascular source for thrombo-embolism by echocardiography, ECG, and vascular sonography demands additional vascular imaging that allows an entire vascular assessment to detect rare sources for embolism.

Panel A. Three-dimensional volume rendering computed tomography image. Aneurysm of the left posterior circumflex humeral artery (white arrow) and thromboembolic occlusion of the left brachial artery bifurcation (white arrowhead).

Panel B. Transverse computed tomography image. Aneurysm of the left posterior circumflex humeral artery (white arrow).

